

Loprazolam (pron. low-praz-o-lamb)

What is loprazolam used for?

Loprazolam is usually used to help treat insomnia i.e. problems getting to sleep. It is one of a number of medicines called "hypnotics" and is sometimes called a "benzodiazepine". It is available as tablets.

What is the usual dose of loprazolam?

The usual dose of loprazolam is around 1-2mg at bedtime.

How should I take loprazolam?

Swallow the tablets with at least half a glass of water whilst sitting or standing. This is to make sure that they reach the stomach and do not stick in your throat.

When should I take loprazolam?

Take your loprazolam at bedtime, probably about half an hour before you go to bed. There is no problem about taking loprazolam with or after food.

What are the alternatives to loprazolam?

There are many other medicines (e.g. the "Z hypnotics"), talking therapies and treatments for insomnia, and the causes of insomnia. See our "Handy charts" to help you compare the medicines, how they work and their side effects.

How long will loprazolam take to work?

Loprazolam usually starts to work in about an hour. The effect wears off in about 4-6 hours or so.

How long will I need to keep taking loprazolam for?

Please look one of the "Handy charts" for more help and advice on how long you might need to take it for. It is not a good idea to take it every night as the effect may start to wear off.

Is loprazolam addictive?

Loprazolam is not addictive as such, but you can get some symptoms if you stop it suddenly (see the next question). It is also true that it can be abused or misused by some people.

Can I stop taking loprazolam suddenly?

Some people get something called "rebound insomnia" when they stop a hypnotic. This means they find it even more difficult to get to sleep than before. It is best to take loprazolam only when you need it but if you have taken it every night for more than a few weeks, it might be best to reduce the dose slowly over a few days. You could discuss this fully with your doctor, nurse or pharmacist.

What should I do if I forget to take loprazolam?

This is not really a problem.

Can I drink alcohol while I am taking loprazolam?

If you drink alcohol while taking loprazolam it may make you feel more sleepy, especially the next morning. This is particularly important if you need to drive or operate machinery and you must seek advice on this.

Will loprazolam affect my other medication?

Loprazolam has only a few interactions with other medicines.

- The effect of loprazolam can be decreased by caffeine (e.g. in tea, coffee, cola drinks, some energy drinks), theophylline or aminophylline (for asthma)
- If loprazolam is taken with alcohol, it will cause more sleepiness. You will also get more sleepiness if you take loprazolam with anything else that causes sleepiness e.g. some antihistamines, antidepressants, pain killers, antipsychotics, anticonvulsants and many other medicines
- You should have no problems with "The Contraceptive Pill" and loprazolam

Not all of these interactions happen in everyone. Some of these medicines can still be used together but you will need to follow your doctor's instructions carefully. There are many other possible drug interactions.

What sort of side-effects might occur if I am taking loprazolam?

The table below will show you some of the main side effects you might get from loprazolam.

Side effect	What happens	What to do about it
COMMON (<i>more than about 1 in 10 people might get these</i>)		
Fatigue and sleepiness during the day	You feel tired or drowsy, especially in the morning.	If you feel like this for more than a week after starting loprazolam, tell your doctor. It may be possible to reduce your dose.
Dizziness	Feeling light-headed and faint.	Don't stand up too quickly. Try and lie or sit down if you feel it coming on. Don't drive.
UNCOMMON (<i>less than about 1 in 10 people might get these</i>)		
Ataxia	Being unsteady on your feet.	Discuss with your doctor when you next see him or her.
RARE (<i>less than about 1 in 100 people might get these</i>)		
Aggression	Feeling excitable. You may be talkative, unfriendly or disinhibited.	Discuss this with your doctor. He or she may want to adjust your medicine or dose.
Confusion	Your mind is all mixed up or confused.	Discuss with your doctor when you next see him or her. He or she may want to adjust your medicine or dose.
Hypotension	Low blood pressure – this can make you feel dizzy, particularly when you stand up.	It is not dangerous. Don't stand up too quickly. If you feel dizzy, don't drive.
Amnesia	Loss of short-term memory. Difficulty in remembering.	It is not dangerous. Discuss with your doctor if you are worried.
Rashes	Blotches seen anywhere.	Stop taking loprazolam and see your doctor now.
OTHER		
Rebound insomnia	More difficult to get to sleep if you stop taking loprazolam.	If you feel like this for more than a week after stopping loprazolam, tell your doctor.

Do not be worried by this list of side effects. Some people get no side effects at all and others may get some effects that are not listed in this table. If you think you might have a side effect to your medicine, you should ask your doctor, nurse or pharmacist. If you want to know more, go to our website for links to other websites with more information.

Will I need a blood test if I am taking loprazolam?

You should not need a blood test while you are taking loprazolam.

Can I drive while I am taking loprazolam?

You may feel a bit sleepy the next morning after taking loprazolam. You should be careful as it may slow down your reaction times. Until this wears off, or you know how loprazolam affects you, do not drive or operate machinery.

Prepared for you by Choice and Medication

This leaflet is to help you understand about your medicine. It is not a manufacturer's Patient Information Leaflet. You may find lots more on the internet but beware as internet-based information is not always accurate. The "Handy charts" will help you compare the main medicines for each condition, how they work and their side effects.